


Section 6.4 The Gram-Schmidt Process

Goal: Form an orthogonal basis for a subspace W .

EXAMPLE: Suppose $W = \text{Span}\{\mathbf{x}_1, \mathbf{x}_2\}$ where $\mathbf{x}_1 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$ and $\mathbf{x}_2 = \begin{bmatrix} 2 \\ 2 \\ 3 \end{bmatrix}$. Find an orthogonal basis $\{\mathbf{v}_1, \mathbf{v}_2\}$ for W .



Let

$$\mathbf{v}_1 = \mathbf{x}_1 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}.$$

$$\hat{\mathbf{y}} = \text{proj}_{\mathbf{v}_1} \mathbf{x}_2 = \frac{\mathbf{x}_2 \cdot \mathbf{v}_1}{\mathbf{v}_1 \cdot \mathbf{v}_1} \mathbf{v}_1$$

and

$$\mathbf{v}_2 = \mathbf{x}_2 - \hat{\mathbf{y}} = \mathbf{x}_2 - \frac{\mathbf{x}_2 \cdot \mathbf{v}_1}{\mathbf{v}_1 \cdot \mathbf{v}_1} \mathbf{v}_1 = \begin{bmatrix} 2 \\ 2 \\ 3 \end{bmatrix} - \frac{4}{2} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 3 \end{bmatrix}$$

(component of \mathbf{x}_2 orthogonal to \mathbf{x}_1)

EXAMPLE: Suppose $\{\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3\}$ is a basis for a subspace W of \mathbf{R}^4 . Describe an orthogonal basis for W .

Solution: Let

$$\mathbf{v}_1 = \mathbf{x}_1 \text{ and } \mathbf{v}_2 = \mathbf{x}_2 - \frac{\mathbf{x}_2 \cdot \mathbf{v}_1}{\mathbf{v}_1 \cdot \mathbf{v}_1} \mathbf{v}_1.$$

$\{\mathbf{v}_1, \mathbf{v}_2\}$ is an orthogonal basis for $\text{Span}\{\mathbf{x}_1, \mathbf{x}_2\}$.

Let

$$\mathbf{v}_3 = \mathbf{x}_3 - \frac{\mathbf{x}_3 \cdot \mathbf{v}_1}{\mathbf{v}_1 \cdot \mathbf{v}_1} \mathbf{v}_1 - \frac{\mathbf{x}_3 \cdot \mathbf{v}_2}{\mathbf{v}_2 \cdot \mathbf{v}_2} \mathbf{v}_2$$

(component of \mathbf{x}_3 orthogonal to $\text{Span}\{\mathbf{x}_1, \mathbf{x}_2\}$)

Note that \mathbf{v}_3 is in W . Why?

$\Rightarrow \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ is an orthogonal basis for W .

THEOREM 11 THE GRAM-SCHMIDT PROCESS

Given a basis $\{\mathbf{x}_1, \dots, \mathbf{x}_p\}$ for a subspace W of \mathbf{R}^n , define

$$\mathbf{v}_1 = \mathbf{x}_1$$

$$\mathbf{v}_2 = \mathbf{x}_2 - \frac{\mathbf{x}_2 \cdot \mathbf{v}_1}{\mathbf{v}_1 \cdot \mathbf{v}_1} \mathbf{v}_1$$

$$\mathbf{v}_3 = \mathbf{x}_3 - \frac{\mathbf{x}_3 \cdot \mathbf{v}_1}{\mathbf{v}_1 \cdot \mathbf{v}_1} \mathbf{v}_1 - \frac{\mathbf{x}_3 \cdot \mathbf{v}_2}{\mathbf{v}_2 \cdot \mathbf{v}_2} \mathbf{v}_2$$

\vdots

$$\mathbf{v}_p = \mathbf{x}_p - \frac{\mathbf{x}_p \cdot \mathbf{v}_1}{\mathbf{v}_1 \cdot \mathbf{v}_1} \mathbf{v}_1 - \frac{\mathbf{x}_p \cdot \mathbf{v}_2}{\mathbf{v}_2 \cdot \mathbf{v}_2} \mathbf{v}_2 - \dots - \frac{\mathbf{x}_p \cdot \mathbf{v}_{p-1}}{\mathbf{v}_{p-1} \cdot \mathbf{v}_{p-1}} \mathbf{v}_{p-1}$$

Then $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ is an orthogonal basis for W and

$$\text{Span}\{\mathbf{x}_1, \dots, \mathbf{x}_p\} = \text{Span}\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$$

EXAMPLE Suppose $\{\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3\}$, where

$$\mathbf{x}_1 = \begin{bmatrix} 1 \\ 2 \\ 3 \\ 0 \end{bmatrix}, \mathbf{x}_2 = \begin{bmatrix} 1 \\ 2 \\ 0 \\ 0 \end{bmatrix}, \mathbf{x}_3 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 1 \end{bmatrix}, \text{ is a basis for a}$$

subspace W of \mathbf{R}^4 . Describe an orthogonal basis for W .

$$\text{Solution: } \mathbf{v}_1 = \mathbf{x}_1 = \begin{bmatrix} 1 \\ 2 \\ 3 \\ 0 \end{bmatrix} \text{ and}$$

$$\mathbf{v}_2 = \mathbf{x}_2 - \frac{\mathbf{x}_2 \cdot \mathbf{v}_1}{\mathbf{v}_1 \cdot \mathbf{v}_1} \mathbf{v}_1 = \begin{bmatrix} 1 \\ 2 \\ 0 \\ 0 \end{bmatrix} - \frac{5}{14} \begin{bmatrix} 1 \\ 2 \\ 3 \\ 0 \end{bmatrix} = \begin{bmatrix} \frac{9}{14} \\ \frac{9}{7} \\ -\frac{15}{14} \\ 0 \end{bmatrix}$$

$$\text{Replace } \mathbf{v}_2 \text{ with } 14\mathbf{v}_2 : \mathbf{v}_2 = 14 \begin{bmatrix} \frac{9}{14} \\ \frac{9}{7} \\ -\frac{15}{14} \\ 0 \end{bmatrix} = \begin{bmatrix} 9 \\ 18 \\ -15 \\ 0 \end{bmatrix}$$

(optional step - to make \mathbf{v}_2 easier to work with in the next step)

$$\mathbf{v}_3 = \mathbf{x}_3 - \frac{\mathbf{x}_3 \cdot \mathbf{v}_1}{\mathbf{v}_1 \cdot \mathbf{v}_1} \mathbf{v}_1 - \frac{\mathbf{x}_3 \cdot \mathbf{v}_2}{\mathbf{v}_2 \cdot \mathbf{v}_2} \mathbf{v}_2$$

$$\mathbf{v}_3 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 1 \end{bmatrix} - \frac{1}{14} \begin{bmatrix} 1 \\ 2 \\ 3 \\ 0 \end{bmatrix} - \frac{9}{630} \begin{bmatrix} 9 \\ 18 \\ -15 \\ 0 \end{bmatrix}$$

$$= \begin{bmatrix} 1 \\ 0 \\ 0 \\ 1 \end{bmatrix} - \frac{1}{14} \begin{bmatrix} 1 \\ 2 \\ 3 \\ 0 \end{bmatrix} - \frac{1}{70} \begin{bmatrix} 9 \\ 18 \\ -15 \\ 0 \end{bmatrix} = \begin{bmatrix} \frac{4}{5} \\ -\frac{2}{5} \\ 0 \\ 1 \end{bmatrix}$$

$$\text{Rescale (optional): } \mathbf{v}_3 = \begin{bmatrix} 4 \\ -2 \\ 0 \\ 5 \end{bmatrix}$$

Orthogonal Basis for W :

$$\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\} = \left\{ \begin{bmatrix} 1 \\ 2 \\ 3 \\ 0 \end{bmatrix}, \begin{bmatrix} 9 \\ 18 \\ -15 \\ 0 \end{bmatrix}, \begin{bmatrix} 4 \\ -2 \\ 0 \\ 5 \end{bmatrix} \right\}$$

Orthonormal Basis

Suppose the following is an orthogonal basis for subspace

$$W = \text{span} \left\{ \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 2 \\ 3 \end{bmatrix} \right\} :$$

$$\{\mathbf{v}_1, \mathbf{v}_2\} = \left\{ \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 3 \end{bmatrix} \right\}$$

Rescale to form unit vectors:

$$\mathbf{u}_1 = \frac{1}{\|\mathbf{v}_1\|} \mathbf{v}_1 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \\ 0 \end{bmatrix}$$

$$\mathbf{u}_2 = \frac{1}{\|\mathbf{v}_2\|} \mathbf{v}_2 = \frac{1}{3} \begin{bmatrix} 0 \\ 0 \\ 3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

Orthonormal basis for W : $\{\mathbf{u}_1, \mathbf{u}_2\}$

QR Factorization

THEOREM 12 (The QR Factorization)

If A is an $m \times n$ matrix with linearly independent columns, then A can be factored as $A = QR$, where Q is an $m \times n$ matrix whose columns form an orthogonal basis for $\text{Col } A$ and R is an $n \times n$ upper triangular invertible matrix with positive entries on its main diagonal.

EXAMPLE Find the QR factorization of $A = \begin{bmatrix} 1 & 2 \\ 1 & 2 \\ 0 & 3 \end{bmatrix}$.

Solution: Use the Gram Schmidt process to find

$\left\{ \begin{bmatrix} \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$ which is an orthonormal basis for

$\text{col } A = \text{span} \left\{ \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 2 \\ 3 \end{bmatrix} \right\}$. So $Q = \begin{bmatrix} \frac{1}{\sqrt{2}} & 0 \\ \frac{1}{\sqrt{2}} & 0 \\ 0 & 1 \end{bmatrix}$.

Since Q has orthonormal columns, $Q^T Q = I$. So if $A = QR$, then

$$A = QR$$

$$R = Q^T A = \begin{bmatrix} \frac{1}{2}\sqrt{2} & \frac{1}{2}\sqrt{2} & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 1 & 2 \\ 0 & 3 \end{bmatrix} = \begin{bmatrix} \sqrt{2} & 2\sqrt{2} \\ 0 & 3 \end{bmatrix}.$$